

Rhodora

JOURNAL OF THE
NEW ENGLAND BOTANICAL CLUB

Conducted and published for the Club, by

MERRITT LYNDON FERNALD, Editor-in-Chief

ALBERT FREDERICK HILL
STUART KIMBALL HARRIS
RALPH CARLETON BEAN

} Associate Editors

Vol. 52.

April, 1950.

No. 616.

CONTENTS:

A Small Fascicle of Novelties. <i>M. L. Fernald</i>	61
I. A Study of <i>Habenaria orbiculata</i>	61
II. A Southeastern <i>Nuphar</i>	65
III. A Southern <i>Amelanchier</i>	67
IV. The Autumnal Perennial <i>Gentian</i> of the Desert.....	68
V. Three minor Transfers.....	71
<i>Arenaria uniflora</i> . <i>M. L. Fernald and B. G. Schubert</i>	71
Nomenclatural and other Notes on Mosses. <i>Herbert Habeeb</i>	72
<i>Holcus mollis</i> in New Hampshire. <i>Arthur Stanley Pease</i>	75
A New Flora for the San Francisco Bay Area (Review). <i>R. C. Rollins</i>	76

The New England Botanical Club, Inc.

8 and 10 West King St., Lancaster, Pa.

Botanical Museum, Oxford St., Cambridge 38, Mass.

RHODORA.—A monthly journal of botany, devoted primarily to the flora of the Gray's Manual Range and regions floristically related. Price, \$4.00 per year, net, postpaid, in funds payable at par in United States currency in Boston; single copies (if available) of not more than 24 pages and with 1 plate, 40 cents, numbers of more than 24 pages or with more than 1 plate mostly at higher prices (see 3rd cover-page). Back volumes can be supplied at \$4.00. Some single numbers from these volumes can be supplied only at advanced prices (see 3rd cover-page). Somewhat reduced rates for complete sets can be obtained on application to Dr. Hill. Notes and short scientific papers, relating directly or indirectly to the plants of the northeastern states, will be considered for publication to the extent that the limited space of the journal permits. Illustrations can be used only if the cost of engraver's blocks is met through the author or his institution. Forms may be closed five weeks in advance of publication. Authors (of more than two pages of print) will receive 15 copies of the issue in which their contributions appear, if they request them when returning proof. Extracted reprints, if ordered in advance, will be furnished at cost.

Address manuscripts and proofs to

M. L. Fernald, 14 Hawthorn Street, Cambridge 38, Mass.

Subscriptions (making all remittances payable to RHODORA) to Dr. A. F. Hill, 8 W. King St., Lancaster, Pa., or, preferably, Botanical Museum, Oxford St., Cambridge 38, Mass.

Entered as second-class matter March 9, 1929, at the post office at Lancaster, Pa., under the Act of March 3, 1879.

INTELLIGENCER PRINTING COMPANY

Specialists in Scientific and Technical Publications

EIGHT WEST KING ST., LANCASTER, PA.

CONTRIBUTIONS FROM THE GRAY HERBARIUM

No. XXXIV.	Reprinted from Proc. Am. Acad. xxxviii. 77-269. 1902. Flora of the Galapagos Islands. With 3 plates. By B. L. Robinson.	\$1.75
No. XXXV.	Reprinted from Proc. Am. Acad. xxxix. 60-120. 1903. New and otherwise Noteworthy Angiosperms from Mexico and Central America. By J. M. Greenman.	\$.50
No. XXXI.	Reprinted from Proc. Am. Acad. xli. 235-278. 1905. 1. Descriptions of Spermatophytes from the Southwestern United States, Mexico, and Central America. By J. M. Greenman. 2. Diagnoses and Notes Relating to American Eupatorieae. By B. L. Robinson.	\$.40
No. XXXII.	Reprinted from Proc. Am. Acad. xlii. 3-48. 1906. 1. Revision of the Genus Piqueria. 2. Revision of the Genus Ophryosporus. 3. The Genus Helogyne and its Synonyms. 4. Diagnoses and Synonymy of Eupatorieae and of certain other Compositae which have been classed with them. By B. L. Robinson.	\$.60

GRAY HERBARIUM of HARVARD UNIVERSITY, Cambridge 38, Mass.



Digitized by the Internet Archive
in 2023 with funding from
Kahle/Austin Foundation



Photo. B. G. Schubert

HABENARIA ORBICULATA, var. *LEHORSII*: FIG. 1, inflorescence, $\times 1$, from TYPE; FIG. 2, inflorescence, $\times 1$, from TOPOTYPE.

H. ORBICULATA, passing to var. *LEHORSII*: FIG. 3, inflorescence, $\times 1$, from Lookout Mt., Bonne Bay, Newfoundland; FIG. 4, fruit, $\times 1$, from Bay of Islands, Newfoundland.

H. ORBICULATA, var. *MENZIESII*: inflorescence, $\times 1$, from Flathead Lake, Montana.

Rhodora

JOURNAL OF

THE NEW ENGLAND BOTANICAL CLUB

Vol. 52.

April, 1950

No. 616

A SMALL FASCICLE OF NOVELTIES

M. L. FERNALD

(Plates 1157—1160)

IN partially clearing the record before the issue of the 8th edition of Gray's Manual it is important to discuss some interesting plants of our flora which have been held over until they could receive proper study. The first two discussions are accompanied by four of Dr. Schubert's very clear plates. The cost of their reproduction has been generously met by Mr. BAYARD LONG.

I. A STUDY OF HABENARIA ORBICULATA (Plates 1157 and 1158)

HABENARIA ORBICULATA (Pursh) Torr., var. **Lehorsii**, var. nov. (TAB. 1157, FIG. 1 et 2, $\times 1$), planta parva foliis 6–9 cm. longis 4–5 cm. latis; scapo 7–11 cm. alto; racemo 2–5.5 cm. longo, 2–4 cm. crasso floribus confertis, internodiis imis 1–8 mm. longis; pedicellis 2–4 mm. longis crassis; labio 7–10 mm. longo; calcare 0.8–1.5 cm. longo.—ST. PIERRE ET MIQUELON: bare spots among Ericaceous shrubs on a denuded hill, Cap à l'Aigle, July 25, 1945, Mathurin Le Hors (TYPE in Herb. Gray); same station, August 2, 1949, Le Hors.

Var. *Lehorsii* is the most extreme departure from typical *Habenaria orbiculata* (our T. 1158, FIGS. 1 and 2) of moist to dry woodlands from southern and western Newfoundland to Thunder Bay District, Ontario, and Minnesota, thence south to Nova Scotia and the Northern States and on the uplands to South Carolina and Tennessee. In its more northeastern and bleak habitats, the latter (typical) plants somewhat approach var. *Lehorsii*; and some of them, especially on or among the mountains of western Newfoundland (T. 1157, FIGS. 3 and 4, $\times 1$) go as well with that

as with true *H. orbiculata*. The type- and topotype-series of var. *Lehorsii* stand off on several characters from the continental (and rarely insular) plant as these are displayed in the 153 collections of the latter before me. These are summarized below and the details of the inflorescence are shown in T. 1158, FIGS. 1 and 2:

H. ORBICULATA (typical), T. 1158, FIGS. 1 and 2, $\times 1$: Leaves 0.6–2 dm. long, 0.4–2 dm. broad; scape 1.5–5 dm. high, with (1–)2–5 bracts; raceme lax and open, 0.7–2.5 dm. long and 3.5–8 cm. in diameter, the flowers 0.8–2.5 cm. apart; pedicels filiform, 6–13 mm. long in maturity; spur 1.6–2.7 cm. long; lip 1–1.5 cm. long.

Var. *LEHORSII*: Leaves 6–9 cm. long, 4–5.5 cm. broad; scape 7–11 cm. high, with 0–2 (rarely 3) bracts; raceme dense, 2–5.5 cm. long, 2–4 cm. in diameter, the lower flowers 1–8 mm. apart; pedicels stout, 2–4 mm. long; spur 8–15 mm. long; lip 7–10 mm. long.

Habenaria orbiculata, var. *Lehorsii* of open heathy barrens is thus quite parallel with *H. Hookeri* Torr., var. *abbreviata* Fernald in RHODORA, xxxv. 239, t. 252 (1933), that dwarf and endemic Newfoundland variety of a continental woodland species known likewise on open heathy barrens, which are carpeted by *Juniperus communis*, var. *saxatilis* Pallas, *Empetrum nigrum* and *Dryas integrifolia* (l. c. 54). When *H. Hookeri*, var. *abbreviata* was described I pointed out (RHODORA, l. c. 8, 9, 87 and 88) the remarkable occurrence on open barrens or mountain-tablelands of a phase of *H. orbiculata* which greatly puzzled Long, Fogg and myself, saying: "other southern species on French Island [Bay of Islands] which seemed absolutely out of place in close proximity to [dominantly Arctic] *Silene acaulis* and *Saxifraga oppositifolia* (MAP 1) were *Cypripedium reginae* Walt . . . and *Habenaria orbiculata* (MAP 9) . . . *Habenaria orbiculata* was so abundant as to be almost dominant in dry peat and gravel within a stone's throw of the open shore. In New England and eastern Canada we had always thought of it as an orchid of the richest of old woodlands; but its choice of habitat on bleak French Island [fruiting raceme in T. 1157, FIG. 4, $\times 1$] was only a breaking of the ice for our next station for it. In 1929 we found it [flowering raceme as T. 1157, FIG. 3, $\times 1$] vying with the arctic *Scirpus cespitosus* var. *callosus* (MAP 10) for possession of a sphagnum pocket on the high tableland of Lookout Mountain at Bonne Bay". As to the station on Lookout Mountain, at an altitude approaching 500 m., whereas close to sea-level at the western base of the mountain the arctic *Oxyria digyna* abounds, the fol-

lowing record was made (pp. 87, 88): "As already stated, *Habenaria orbiculata* (MAP 9) was here growing in turf of the ubiquitous arctic-alpine *Scirpus cespitosus* var. *callosus* (MAP 10). We had difficulty in believing our own eyes and, at my request, Fogg made detailed drawings of its flowers, lest we might forget how the fresh flowers appeared, but we cannot make it anything but *H. orbiculata* of 'rich woods' on the continent".

In the light of the present study it has become evident that the plants of open barrens of western Newfoundland (FIGS. 3 and 4), although with longer and slightly less crowded racemes and larger leaves than in the LeHors material, have the short and thick pedicels of the latter and relatively short spur and lip. They are transitional in all characters.

In summarizing (l. c. 240) in 1933, I said: "I am unable to find any tangible character to separate this extreme series from the larger Alleghenian plant and particularly from the plants of western North America", the latter plant described as *Platanthera Menziesii* Lindl. Gen. Sp. Orchid. Pl. 286 (1935). This plant, however, seems sufficiently definite (our t. 1157, FIG. 5) as to stand as an isolated northwestern variety, rather closer to the Newfoundland plant than to the wide-ranging eastern continental *H. orbiculata*, var. *typica*:

HABENARIA ORBICULATA (Pursh) Torr., var. **Menziesii** (Lindl.), stat. nov. *Platanthera Menziesii* Lindl. Gen. Sp. Orchid. Pl. 286 (1835). *Lysias Menziesii* (Lindl.) Rydb. Fl. Rocky Mts. 178 (1917).

When var. *Menziesii* is separated from the more eastern *Habenaria orbiculata* and *H. macrophylla* Goldie in Edinb. Phil. Journ. vi. 331 (1822), with both of which it is often confused, it stands out, at least as shown in the 12 numbers in the Gray Herbarium, through its relatively small leaves (5–15 cm. long and 4–12 cm. broad; in typical *H. orbiculata* up to 2 dm. long and broad); scape 1–2 dm. high (in typical *H. orbiculata* 1.5–5 dm. high); stricter raceme, the longest scarcely 2 dm. long and with 12–20 remote flowers (longest racemes of eastern typical *H. orbiculata* up to 2.5 dm. long and with 25–50 flowers); pedicels stoutish (as opposed to filiform); spur at most 2.2 cm. long (in the East up to 2.7 cm.). Although the thickish pedicels usually found in var. *Menziesii* had earlier suggested to me those of var.

Lehorsii and transitions to it, the shorter pedicels, crowded flowers and shorter spur and lip clearly distinguish the latter from the former.

Much confusion has arisen through the inclusion in *Habenaria orbiculata* or *Lysias orbiculata* (Pursh) Rydb., originally described by Pursh as *Orchis orbiculata* Pursh. Fl. Am. Sept. ii. 588 (1814) from "shady beech-woods: on the mountains of Pennsylvania and Virginia", of the technically quite different *H. macrophylla* Goldie, described from Montreal. There can be no doubt as to the identity of *H. macrophylla*. Goldie compared it with *H. orbiculata*, its leaves said to be "four times as large as those of *H. orbiculata*, measuring from six to eight inches in length, very thin and pellucid . . . Flowers large, white, . . . and arranged in a lax spike of about five or six inches in length . . . *labellum* . . . linear, as long as the germen . . . *Germen* about an inch in length." A very illuminating review of early misidentifications of *H. macrophylla* and *H. orbiculata*, with beautiful drawings of the flowers of the two, was published by Ames in RHODORA, viii. 1-5 with figs. (1906). After his most careful discussion Ames gave the measurements of the spurs of *H. orbiculata* (ranging from 16 to 27, most often 18-20 mm. long) and of *H. macrophylla* (ranging from 32 to 43, most often 35-37 mm. long); while the type of "*Platanthera Menziesii* is the short-spurred form of *H. orbiculata*". With such a careful study at hand, it is unfortunate that *H. macrophylla* (our t. 1158, figs. 3 and 4, $\times 1$) and *H. orbiculata* (our t. 1158, figs. 1 and 2, $\times 1$) should still be confused. The inclusion of *H. macrophylla* by Abrams in his Ill. Fl. Pacific States, i. 477 (1940) in the synonymy of the carefully illustrated *Lysias orbiculata* (his fig. 1168) may account for his "Scape short, 3-6 dm. high". At least, the material of the northwestern plant illustrated by him (our *H. orbiculata*, var. *Menziesii*) in the Gray Herbarium shows scapes only 1-2 dm. high, while the scape of typical *H. orbiculata* rarely reaches a height of 5 dm.¹

Although *Habenaria macrophylla* and typical *H. orbiculata* are both woodland plants, they rarely, if ever, grow together. In the Gray Herbarium and that of the New England Botanical Club there are 135 collections from eastern Canada (Quebec

¹ Through the dates of publication given by Abrams (1826 for Pursh's basic *Orchis rotundifolia*, 1822 for Goldie's *Habenaria macrophylla*) one would infer that the latter name has priority. By correcting the Pursh date to 1814 this difficulty is removed.



Photo. B. G. Schubert

HABENARIA ORBICULATA (var. TYPICA): FIG. 1, flowering inflorescence, $\times 1$, from New Hampshire; FIG. 2, fruit, $\times 1$, from New Hampshire.

H. MACROPHYLLA: FIG. 3, unusually small inflorescence, $\times 1$, from Vermont; FIG. 4, fruit, $\times 1$, from Maine.

eastward and southward) and northern New England of these two species. A listing of all localities shows only 7 cases in which both species have been collected in the same township or similar large area. This is well illustrated in Newfoundland, where variations of *H. orbiculata* occur on the western side of the Island and near the southern coast west of the Avalon Peninsula; but only *H. macrophylla* is known from the Avalon Peninsula. Furthermore, in general *H. macrophylla* occurs more often in richer soil than does *H. orbiculata*: such statements of habitat as "woods" or "damp" or "dry woods" are not distinctive but many labels of *H. macrophylla* indicate "deep rich woods", "hardwoods", "rich deciduous woods", "old mixed woods", "old maple-grove", etc., specifications rarely found on labels of *H. orbiculata*. As supplemental to the characters usually recognized as distinguishing *H. macrophylla*, I find that its full-grown capsules (our FIG. 4) range from 1.8–2.5 cm. long, with a thickness of 6–8 mm.; those of *H. orbiculata* (FIG. 2) 1.3–2 cm. long and 4–6 mm. thick.

EXPLANATION OF PLATES

PLATE 1157, all figs. $\times 1$. FIGS. 1 and 2, racemes of *HABENARIA ORBICULATA* (Pursh) Torr., var. *LEHORSH* Fernald: FIG. 1 from TYPE; FIG. 2 from TOPOTYPE. FIGS. 3 and 4, *H. ORBICULATA* passing into var. *LEHORSH*: FIG. 3 from turf of *Scirpus caespitosus* L., var. *callosus* Bigel. and in *Sphagnum*, wet peaty bog-barrens at 400–550 m. alt., tableland of Lookout Mountain, Bonne Bay, Newfoundland, Fernald, Long & Fogg, no. 1554; FIG. 4, fruit, sphagnous boggy marsh, French (or Tweed) Island, Bay of Islands, Newfoundland, Fernald, Long & Fogg, no. 196. FIG. 5, raceme of *H. ORBICULATA*, var. *MENZIESII* (Lindl.) Fernald from Big Fork, Flathead Lake and vicinity, Montana, July 15, 1908, Mrs. Joseph Clemens.

PLATE 1158, all figs. $\times 1$. FIGS. 1 and 2, portions of racemes of *HABENARIA ORBICULATA*, var. *TYPICA*: FIG. 1, flowers from Whitefield, New Hampshire, July 28, 1895, Walter Deane; FIG. 2, fruits from Franconia, New Hampshire, September 24, 1895, E. & C. E. Faxon. FIGS. 3 and 4, *H. MACROPHYLLA* Goldie: FIG. 3, unusually small raceme from Willoughby, Vermont, July 14, 1898, G. G. Kennedy; FIG. 4, fruits, Douglas Hill, Sebago, Maine, Fernald, Long & Norton, no. 13,318.

II. A SOUTHEASTERN NUPHAR (Plates 1159 and 1160)

NUPHAR puteorum, n. sp.¹ TAB. 1159 et TAB. 1160, FIG. 1–6. Similar to *N. ovatum* (Miller & Standley) Standley, but leaf-blades usually more tapering to apex, sparsely and minutely silky-puberulent beneath, the sinus 2–4 cm. broad near the middle, petiole subterete, densely puberulent; fruit 2.5–3.8 cm.

¹ *NUPHAR puteorum*, sp. nov., a *N. ovato* differt foliis ad apicem plus ovato-attenuatis, sinu latiore (medio 2–4 cm. lato); fructibus 2.3–3.5 cm. diametro; disco 1.3–1.5 cm. diametro, margine valde undulato, centro plerumque papillifero; seminibus quadrato-obovatis 4–5 mm. longis dorso puberulis.

high, 2.3–3.5 cm. thick, obscurely ribbed; rim of disk undulate, 1.3–1.5 cm. in diameter; disk deeply depressed, the center usually papillate; rays extending nearly to margin of disk; seeds quadrate-obovate, 4–5 mm. high, puberulent, at least on the back.—Sussex County, VIRGINIA: deep muddy pit at margin of Chappell's Mill-pond (Honey Pond), west of Lumberton, July 3, 1942, *Fernald & Long*, no. 14,326 (TYPE in Herb. Gray.; ISOTYPE in Herb. Phil. Acad.).

Nuphar puteorum occupied the center of a deep pit dug at the margin of Chappell's (or Honey) Pond. The very slippery and muddy border of the pit was so treacherous that we were able, with the aid of forked branches of trees, to pull ashore only a few broken-off flowering and fruiting stalks and some characteristic leaves. So treacherous was the footing and so indefinite the possible depth of the pit that we planned to return with a strong rope, as a safeguard, and secure better material, a responsibility now left to others. Hence the specific name, from *puteus*, a pit (not from *puteo*, to stink, although, after our half-drowning and unsavory experience from marsh-fumes in getting it, we inclined toward the latter source of the specific name).

Nuphar puteorum apparently belongs in the extreme southern and southwestern series of species with the lower leaf-surface more or less puberulent; and, judging from the monograph by Gerrit S. Miller & Paul C. Standley, *North American Species of Nymphaea*, Contrib. U. S. Nat. Herb. xvi³. (1912), it is nearest allied to the Texan *Nuphar ovatum* (Mill. & Standl.) Standl. From that monograph some details (our T. 1160, FIGS. 7 and 8) are here reproduced. The leaf-blade of *N. puteorum* is more ovate than that of *N. ovatum* and its sinus broader, 2–4 cm. across at the middle, the narrow sinus of *N. ovatum* shown as only one-tenth that breadth. The fruit (FIG. 7) of the latter species is 3–4 cm. in diameter and with an entire summit-rim 2.8–4 cm. across; whereas the fruit (T. 1159, FIG. 2) of the Virginian plant is 2.2–3.5 cm. in diameter and with the strongly scalloped or undulate rim only 1.3–1.5 cm. across. The depressed stigmatic disk (T. 1160, FIG. 8) of the Texan plant is entire, 2.2–3 cm. across and with a smooth center, its rays failing by 2–4 mm. to reach the margin; the strongly undulate-margined disk (FIGS. 3 and 4, $\times 3$) of *N. puteorum* is at most 1.5 cm. in diameter and its center often bears 1 to several papillae which resemble short stigmatic rays; and the rays extend




Photo. B. G. Schubert

NUPHAR PUTEORUM (all figs. from TYPE): FIG. 1, leaf, flower and fruit, $\times$ ca. $\frac{1}{3}$; FIG. 2, cross-section of petiole, $\times$ 1; FIG. 3, surface of petiole, to show pubescence, $\times$ 5; FIG. 4, lower surface of leaf, with minute pubescence, $\times$ 5.

nearly to the margin of the disk. The seeds (FIG. 9) of *N. ovatum* are 3.5–4 mm. long, 3 mm. in diameter, ovoid to obovoid and lustrously glabrous; those of the Virginian plant (FIG. 5) strongly quadrate-obovate, 4–5 mm. high and puberulent and dull at least on the back (FIG. 6, $\times 5$), a character which is apparently unique in the genus.

The only other species with which *Nuphar puteorum* might be confused is *N. variegatum* Engelm., the wide-ranging northern species also with floating leaves. But, although the petioles and peduncles may be puberulent, they are conspicuously flattened and the lower leaf-surface is glabrous. Its capsules are corrugated above and with nearly entire summit-rim, and its seeds are relatively slender and lustrous. The newly defined species is a quite different plant and is to be expected farther south, especially near the inner margin of the Coastal Plain.

EXPLANATION OF PLATES 1159 and 1160

PLATE 1159. *NUPHAR PUTEORUM*, all figs from the TYPE: FIG. 1, leaf, flower and fruit, $\times$ ca. $\frac{1}{2}$; FIG. 2, cross-section of petiole, $\times$ 1; FIG. 3, surface of petiole, to show pubescence, $\times$ 5; FIG. 4, lower surface of leaf, with minute pubescence, $\times$ 5.

PLATE 1160. FIGS. 1–6, from TYPE of *NUPHAR PUTEORUM*: FIG. 1, flower laid open, $\times$ 1; FIG. 2, capsule, $\times$ 1; FIGS. 3 and 4, disks, to show rays, center and undulate margin, $\times$ 3; FIG. 5, opaque seeds, $\times$ 1; FIG. 6, seeds, showing felted surface, $\times$ 5. FIGS. 7–9, *N. OVATUM* (Mill. & Standl.) Standl.: FIG. 7, larger of the fruits shown by Miller & Standley, $\times$ 1; FIG. 8, diagram of disk and rays, $\times$ 1, after Miller & Standley; FIG. 9, lustrous and glabrous seeds, $\times$ 1, from Driscoll Lake, Camp Fannin, 8 miles northeast of Tyler, Smith County, Texas, *H. E. Moore*, no. 993, as *N. advena* (Ait.) Ait. f.

III. A SOUTHERN AMELANCHIER

AMELANCHIER CANADENSIS (L.) Medic., var. **subintegra**, var. nov., frutex fastigiatus ad 3 vel 4 m. altus; foliis oblongo-ellipticis deinde subcoriaceis integris vel subintegris.—Damp sandy pine-barrens, southeastern VIRGINIA: south of Zuni, Isle of Wight County, July 4, 1942, *Fernald & Long*, no. 14,338; south of Lee's Mill, Isle of Wight Co., October 16, 1941, *Fernald & Long*, no. 13,949, and April 21, 1942 (in anthesis), *Fernald, Long & Abbe*, no. 14,177; east of Cox Landing, south of South Quay, Nansemond County, July 26 and 28, 1939, *Fernald & Long*, no. 10,670; northeast and east of Cox Landing, June 8, 1940, *Fernald & Long*, no. 12,095 (TYPE in Herb. Gray.; ISOTYPE in Herb. Phil. Acad.).

Throughout the pine-barrens of Isle of Wight and Nansemond Counties var. *subintegra* is a characteristic tall fastigate shrub of the wet depressions and damp thickets. In flower and foliage it is like typical and wide-ranging *Amelanchier canadensis* (rang-

ing from Georgia to southeastern and central Maine, central New Hampshire, southwestern Quebec and central-western New York. Typical *A. canadensis*, however, has the margins of the leaves regularly serrate nearly to base, with 6-11 sharp teeth, with acute sinuses, per cm. of margin. Var. *subintegra* at once attracts notice because of the quite entire or only apically slightly toothed leaves. It is consistent in the depressions of the pine-barrens throughout its area and doubtless crosses into eastern North Carolina. In all flowering material seen the young fruits are blasted; consequently no mature fruit has yet been found.

IV. THE AUTUMNAL PERENNIAL GENTIAN OF THE DESERT

In his scholarly *Plants of Southern New Jersey*, the late Dr. Witmer Stone wrote of the very distinct Pine-barren Gentian, the plant which has long passed as *Gentiana porphyrio* J. F. Gmelin, Syst. ii. 462 (1791):

It is probably a matter of individual preference to determine which gentian is the handsomest. The present species, found only in the remote sections of the Pine Barrens, is certainly the least known and to my mind as handsome as any. Its flaring mouth, the delicate markings within, and the intensity of the blue, make it one of the choicest blooms of the region.

It was apparently first discovered by William Bartram, who sent a drawing of it to Edwards, the British naturalist, who published it in his *Gleanings of Natural History*, vol. V., p. 98, 1758, as the "Autumnal Perennial Gentian of the Desert," but it was not properly named until 1791, and then from the Pine Barrens of South Carolina.—Stone, l. c. 640, 641 (1912).

The plate of William Bartram's plant and his brief description of it are unequivocal. The plate, in color, contained, in the style of Mark Catesby, a bird and a plant, these called on p. 97, under Edwards's Chapter xlv, Plate 255, "*The Yellow-rumped Fly-catcher and The Gentian of the Desert*", although beneath the plate itself Edwards's inscription reads "*The Autumnal Gentian of the Desert*". As stated by Stone, Edwards had the plant from William Bartram and he showed it primarily as an attractive perch for the fly-catcher, saying (p. 98):

I received this bird, preserved dry, from Mr. William Bartram, of Pennsylvania, with a drawing of the Gentian only, which I have added to it rather as a decoration, than a part of this *Natural History*; but nevertheless I shall give Mr. Bartram's account of it. He calls it the Autumnal Perennial Gentian of the Desert, and says, that it produces three or four stalks from one root, each of about a foot high, and some stalks produce two flowers: the flowers are of a fine blue colour: the

stalks and leaves green: the flowers keep long in their beauty, and the roots live many years: they are scarce in Pennsylvania. Catesby, in his History of Carolina, vol. I. pag. 70, has given a different species of the American Gentian.

Bartram's description is quite as satisfactory as are many of Catesby or other pre-Linnaean and early post-Linnaean or even Linnaean ones, but until very recently it was not realized that Linnaeus in 1776 gave a name to the plant of Bartram, this antedating by 15 years the name given by Gmelin. The matter is clearly stated by Dr. W. L. McAtee who, in March, 1949, privately issued a 1-page announcement under the title, A LINNAEAN PAPER NEEDING FURTHER ATTENTION. His first paragraph reads:

Near the end of his career, at the request of a London publisher (Jacob Robson), Linnaeus prepared (or perhaps merely added Latin names to a base provided him) "A Catalogue of the Birds, Beasts, Fishes, Insects, Plants & contained in Edwards's Natural History, in Seven Volumes, with their Latin Names." This was printed by Robson in London, 1776, as a 15-page, crown-folio pamphlet in conjunction with "Some Memoirs of the Life and Works of George Edwards" (38 pp.), apparently of his own composition. Probably this "Catalogue" was also separately distributed and from a closing, "N.B.", it is evident that it was prepared as a supplement to Edwards's work, instructions being given as to placing its parts for binding.

The only other *Gentiana autumnalis* is an illegitimate one, although that fact is not brought out in *Index Kewensis*, and the Linnaean name has priority of nearly a century. I refer to *Gentiana autumnalis* Preyer & Zirkel ex Babington in Journ. Linn. Soc. xi. 318 (1871). The latter name was quite illegitimate, since, under the maintained *G. amarella* L., Babington simply remarked: "This is probably the *G. autumnalis* of P. & Z."

The essential bibliography is as follows:

GENTIANA AUTUMNALIS L. Cat. Edwards's Nat. Hist. 11 (1776), the name based on the plate and William Bartram's quoted description in Edwards, Gleanings Nat. Hist. v. 97, 98, t. 255 (1758). *G. angustifolia* Michx. Fl. Bor.-Am. i. 177 (1803), not Vill. (1787). *G. porphyrio* sensu authors, apparently not the restricted (as described) *G. porphyrio* of J. F. Gmel. (see below). *G. Stoneana* Fern. in RHODORA, xli. 555, t. 579 (1939). *G. porphyrio*, forma *Stoneana* (Fern.) Fern. in RHODORA, xliv. 152 (1942).

Forma **albocaerulea** (Fern.), comb. nov. *G. porphyrio*, forma *albocaerulea* Fern. in RHODORA, xliv. 152 (1942).

Forma **albescens** (Fern.), comb. nov. *G. porphyrio*, forma *albescens* Fern. l. c. (1942).

Forma **porphyrio** (J. F. Gmel.), stat. nov. *G. purpurea* Walt. Fl. Carol. 109 (1788), not L. (1753). *G. porphyrio* J. F. Gmel. Syst. ii. 462 (1791).

Gentiana purpurea Walt. was described under the heading, "Corollis infundibuliformibus" in the following words:

purpurea 3. corollis intus laete purpureis; caule
simplici, fol. linearibus

On account of the earlier *Gentiana purpurea*, J. F. Gmelin renamed Walter's plant, without knowing more than the original description, for the purple gallinules (*Porphyrio*), without a word of new description. In 1939, familiar with the intense blue of the corolla of the plant passing as *G. porphyrio* and well-knowing the truly purple color-variations of other southeastern gentians, I felt that, without any extant material of Walter's *G. purpurea*, there was very serious doubt about its identity. I, therefore, named the Pine-barren Gentian with intensely azure corolla and the tube often brown-mottled within *G. Stoneana*. In RHODORA, xliv. 152 (1942), however, accepting the statement of Dr. R. T. Clausen in Bull. Torr. Bot. Cl. lxxviii. 662 (1941), that he had seen "plants" of the "pink phase" which "came originally from near Wilmington, North Carolina", where found by Mrs. J. Norman Henry; and, having seen a specimen with "lavender" flower from New Jersey, I took up the specific name *G. porphyrio*. Somewhat later, in RHODORA, l. c. 238 (1942), quoting the discoverer, I pointed out that the "plants" found by her "after a long and careful search" were a single individual which she removed to her garden and subsequently divided. Bartram originally said "the flowers are of a fine blue colour"; Michaux described his *Gentiana angustifolia* with "Corolla magna, coerulea"; Elliott (Sk. i. 344) said "bright azure"; Chapman (Fl. 356) said "bright blue"; Small (Fl. 931) said "deep or brilliant azure-blue"; Stone (Pl. So. N. J. 640) speaks of the "intensity of the blue"; and Mrs. Henry, in the letter above quoted, said "In 1938 I ran across a large area that was dotted liberally with *Gentiana porphyrio*—the common blue type. As is my custom on finding a large group of any plant, I began to search for an albino. I had no luck . . . but I did find the pink . . . A long and careful search




Photo. B. G. Schubert

NUPHAR PUTEORUM (all figs. from type): FIG. 1, flower laid open, $\times 1$; FIG. 2, capsule, $\times 1$; FIGS. 3 and 4, disks, $\times 3$; FIG. 5, opaque seeds, $\times 1$; FIG. 6, seeds, showing felted surface, $\times 5$.

N. OVATUM: FIG. 7, fruit, $\times 1$, after Miller & Standley; FIG. 8, diagram of disk, $\times 1$, after Miller & Standley; FIG. 9, lustrous and glabrous seeds, $\times 1$, from Smith County, Texas.

revealed that this was the only one of its color. There were no intermediate shades. . . . It has been growing in a pot in a cold frame ever since. This spring [1942] . . . I ventured to divide it in two". To this I added: "It is evident that I was in error in referring to Mrs. Henry's having found 'plants' with pink flowers". It is further evident, if Walter's *Gentiana purpurea* were accurately named and described, that he had the remarkable experience of finding a very rare color-form, without giving evidence that he saw the usual azure-flowered plant.

At any rate, the proper naming of the latter plant from a colored plate and sufficient description of it as *Gentiana autumnalis* L. (1776) allows us to use for it a wholly appropriate name.

V. THREE MINOR TRANSFERS

In order to make them available for use the following transfers are necessary.

ERIANTHUS GIGANTEUS (Walt.) Muhl., var. **compactus** (Nash), comb. nov. *E. compactus* Nash in Bull. Torr. Bot. Cl. xxii. 419 (1895). *E. saccharoides* Michx., var. *compactus* (Nash) Fern. in RHODORA, xlv. 252 (1943).

MALAXIS BRACHYPODA (Gray) Fernald, forma **bifolia** (Mousley), comb. nov. *M. monophyllos* (L.) Sw., forma *bifolia* Mousley in Orchid Rev. xxxv. 356, fig. (1927).

RUBUS IDAEUS L., var. **caudatus** (Robinson & Schrenk), comb. nov. *R. strigosus* Michx., var. *caudatus* Robinson & Schrenk in Can. Rec. Sci. vii. 14 (1896).

ARENARIA UNIFLORA.—In RHODORA, l. 195–197, t. 1103 (1948), we showed that *Stellaria uniflora* Walt. Fl. Carol. 141 (1788) had been generally misinterpreted and that Walter's specimen as well as diagnosis showed it to be identical with *Arenaria brevifolia* Nutt. ex Torr. & Gray, Fl. N. Am. i. 180 (1838).

On account of the different *A. uniflora* Luce (1823) we decided that Nuttall's name must stand under *Arenaria*. We are indebted to Dr. Bassett Maguire for calling to our attention the fact that Muhlenberg, Cat. 45 (1813) properly made the transfer of Walter's binomial to *Arenaria*. Muhlenberg gave (perhaps fortunately, since Walter's species has been so generally misinterpreted) no discussion, merely, under *Arenaria*.

alb. 8. uniflora
 Stellaria Walt. } one-flowered

It is, therefore, evident that we must replace *Arenaria brevifolia* Nutt. ex Torr. & Gray (1838) by *A. uniflora* (Walt.) Muhl. (1813).—M. L. FERNALD AND B. G. SCHUBERT.

NOMENCLATURAL AND OTHER NOTES ON MOSSES

HERBERT HABEEB

FOR a number of years the writer has been collecting and studying North American mosses. Although the mosses of the region have received an excellent and exhaustive treatment in Dr. A. J. Grout's *Moss Flora of North America North of Mexico*, some problems still exist; and in the following discussion the author hopes to clarify a few, or at least to bring them to the consideration of others.

PYLAISIA Bry. Eur. fasc. 46-47 (1851).

Pylaisia Bry. Eur. is a later homonym of the orthographic variant *Pilaisaea* Desvaux, Jour. Bot. 5: 24 (1814) (1813?). It would, therefore, seem desirable to include it on the list of *nomina generica conservanda proposita* for the Musci.

PYLAISIA POLYANTHA Bry. Eur., var. **Jamesii** (Sull. & Lesq.), comb. nov. *Pylaisia Jamesii* Sull. & Lesq. *Schedae ad Musci* Bor. Am. Ed. 2: 63 (1865). *Pylaisiella polyantha Jamesii* (Sull.) E. G. B. Bull. Torrey Club 23: 230 (1896).

I agree with Mrs. E. G. Britton's allocation of a varietal status to *P. Jamesii*, I find that the chief distinguishing character, number of quadrate alar cells present on the leaves, proves to be a variable one, sometimes on leaves from the same plant. In our locality plants closest to *P. polyantha*, having leaves with relatively few quadrate alar cells, usually grow on rock-ledge; while plants of the var. *Jamesii*, having numerous quadrate alar cells on the leaves, grow on trees.

THELIA ASPRELLA Sull., var. **Lescurii** (Sull.), stat. nov. *Thelia Lescurii* Sull. Mosses U. S. 60 (1856); Sull. in Sull. & Lesq. *Schedae ad Musci* Bor. Am. Ed. 1: 54 (1856).

Here the differences, namely relative degree of marginal ciliation in the leaves and the habitat, separating *Thelia asprella*

from *Thelia Lescurii* are not, in my opinion, of specific rank. In this case it would seem that the degree of marginal ciliation in the leaves is associated with the habitat.

DICRANUM SCOPARIUM Hedw., forma **alatum** (Barnes apud Röhl), stat. et comb. nov. *Dicranum Bonjeani* DeNot., var. *alatum* Barnes apud Röhl, Bot. Centralb. **44**: 386 (1890); *Separat-Abdruck* 2 (1890). *D. alatum* Grout, Moss Flora N. Am. **1**: 88 (1937). *D. scoparium* approaching *Bonjeani* or *D. Bonjeani* approaching *scoparium* on herbarium specimens, various authors.

The plants here placed occur much more commonly in the Northeast than either typical *D. scoparium* or typical *D. Bonjeani*. *D. Bonjeani* being admittedly a subspecies (whatever that is in the study of *Musci*) of *D. scoparium*, I see no reason why the intermediate form should be kept under *D. Bonjeani*.

FISSIDENS ADIANTHOIDES Hedw., forma **immarginatus** (Lindb. apud Lesq. & James), stat. nov. *Fissidens adiantoides* Hedw., var. *immarginatus* Lindb. apud Lesq. & James, Man. Mosses N. Am. 88 (1884). *Fissidens adiantoides* Hedw., (var. *immarginatus*) Barnes, Bot. Gaz. **12**: 27 (1887).

Though the character of no perceptible border even at times does not hold for all of the leaves on an individual plant, the student will find it occasionally convenient to name a specimen forma *immarginatus*. Hedwig, in the Species Muscorum, 1801, the starting point for nomenclature in the *Musci*, spells the specific name "*adianthoides*" not "*adiantoides*".

CAMPTOTHECIUM NUTTALLII (Wils.) Bry. Eur. ex Macoun, Cat. Can. Plants **6**: 188 (1892).

Students wishing to use the name, *Camptothecium Nuttallii*, will have to cite it as shown or in some other form, for the combination was not made in the Bryologia Europaea. It was only indicated by a reference of the name, *Hypnum Nuttallii* Wils., to the new genus, *Camptothecium*. Sullivant, Lesquereux and James, and others made the combination, but only in synonymy. Their combinations are not valid according to article 40 of the International Rules of Botanical Nomenclature. Although Grout, in the Moss Flora of North America, wrote *Homalothecium Nuttallii* (Wils.) Grout, the combination was made earlier and should be given as *Homalothecium Nuttallii* (Wils.) Jaeger & Sauerbeck, *Adumbratio* **2**: 376 (1876-77).

PLAGIOPUS OEDERI (Brid.) Limpr., forma **alpinus** (Schwaegr.), stat. et comb. nov. *Bartramia Oederi*, β . var. *alpina* Schwaegr. Species Muscorum Suppl. 1: 2: 50 (1816). *B. Oederi*, β . var. *condensata* Brid. Musc. Rec. Suppl. 3: 87 (1817). *Plagiopus serratus* Brid. Bry. Univ. 1: 596 (1826). *Plagiopus Oederi*, var. *condensatus* Limpr. Deutschl. Laubm. 2: 550 (1893).

This form, condensed by a xerophytic habitat, has been collected by the writer from the canyon at Grand Falls, New Brunswick.

This is also an amusing case. For Bridel, in his *Bryologia Universa*, described as a new genus and species, *Plagiopus serratus* and yet maintained *Bartramia Oederi*, var. *condensata* in this same publication. Later authors showed the former to be synonymous with the latter (see Paris, Index Bryologicus Ed. 1). Recent authors segregate from *Bartramia* the species *Plagiopus Oederi* (Brid.) Limpr. And, Grout, in the Moss Flora of North America, gives *Plagiopus serratus* Brid. as the type-species of the genus, *Plagiopus*. So, we end up with a synonym acting as a type-species. I, therefore, propose that *Plagiopus Oederi* (Brid.) Limpr. be taken up as the type-species of the genus, *Plagiopus*.

DITRICHUM FLEXICAULE (Schwaegr.) Hampe, forma **Estellae**, forma nova. Plantae parvulae caespitosae; caulibus superne flagelliformis cum foliis minimis.—Plants small, crowded; upper stem-parts forming long, slender, brittle, flagellae with reduced leaves.—NEW BRUNSWICK: Grand Falls, ledges in canyon, *Habeeb* 286, May 5, 1944.

This is an extremely reduced form growing in exposed ledge-cracks that are flooded at high water. The plants each year push out new flagellae which are in turn swept away.

GRIMMIA APOCARPA Hedw., var. **ALPICOLA** (Hedw.) Hartm., forma **papillosa** (Jones apud Grout), comb. nov. *Grimmia alpicola* Hedw., var. *rivularis* (Brid.) Broth., forma *papillosa* Jones apud Grout, Moss Flora N. Am. 2: 14 (1933).

I must say that I agree with Dr. Geneva Sayre's views in her discussion of *Grimmia alpicola* Hedw. in the *Bryologist* 49: 3-7 (1946). Here, she supports a varietal rank for *G. alpicola*. Yet, this leaves forma *papillosa* "in the air", so to speak—hence the above combination. From specimens named by Dr. Grout and the description itself, it is hard to see why in the first place forma *papillosa* was under var. *rivularis*, as it is more closely

related to typical *G. alpicola*, from which it differs only in the extreme papillation of the costa and leaf towards the leaf-apex.

GRIMMIA WRIGHTII (Sull.) Aust., var. **Rauei** (Aust.), stat. nov. *Grimmia (coscinodon) Rauei* Aust. Bull. Torrey Club **6**: 46 (1875). *Coscinodon Ravi* (Aust.) Lesq. & James, Man. Mosses N. Am. 155 (1884). C
Lenata

The chief difference between the species and variety is the erose denticulation of the upper leaf-margin found in *G. Wrightii* and margins nearly entire in *G. Rauei*. Again a variable character.

GRIMMIA DONIANA Sm. Flora Brit. **3**: 1198 (1804).

By such specimens, descriptions and plates (see the Bryologia Europaea plates 249, 250, 251) as have been seen, and allowing for a little variation in capsule-lids and basal leaf-cells, I cannot separate *Grimmia montana* Bry. Eur. and *Grimmia alpestris* Nees from *Grimmia Doniana*. It could be that I lack the *grim* judgment of a *Grimmia* specialist, as I fail to see much difference between a calyptra of several near equal slits, said to be mitri-form, and a calyptra of several slits with one slit longer than the others, said to be cucullate.

GRAND FALLS, NEW BRUNSWICK

HOLCUS MOLLIS IN NEW HAMPSHIRE.—An unfamiliar grass collected by me on 19 July, 1949, in low ground by the Maine Central Railroad in Columbia, N. H., precisely matches herbarium specimens and the description in A. S. Hitchcock's *Manual of the Grasses of the United States* (1935), 301 (and fig. 595), of *Holcus mollis* L. (*Ginannia mollis* (L.) Bubani), a species recently adventive from Europe. Hitchcock records its occurrences in America as reaching from Washington to California, with stations also reported from Lewis County, N. Y., and on ballast at Camden, N. J. The Gray Herbarium and the herbarium of the New England Botanical Club show no previously collected specimens from the Eastern United States. This species has panicles rather smaller than those of *H. lanatus*, and the awn of its second floret is exserted and geniculate, instead of hook-like as in *H. lanatus*.—ARTHUR STANLEY PEASE, Harvard University.

A NEW FLORA FOR THE SAN FRANCISCO BAY AREA¹—One of the delights and compensations of the systematic botanist is his enforced intimate contacts with living dynamic populations of plants while studying them in their wild settings. In *Marin Flora*, Howell conveys to the reader a certain sense of the enjoyment he must have experienced while tramping the Marin hills gathering first hand information about the plants growing there. This book is an account of the plants growing in a single county comprising 529 square miles, but it should be usable in a much larger area of central coastal California. It is based upon an intensive study carried on for more than a ten-year period by its author. However, as is usual in systematics, it reflects the work of several contemporary botanists and many botanists of preceding generations.

Following a description of the physiographic features of Marin County, a section of the book is devoted to "life zones and plant associations". Here the thoughtful student may gain some precise information about the kinds of plants found growing together in various areas and habitats of the region covered. Also included are twenty-four remarkably clear photographs taken by Charles T. Townsend of a variety of views and plants. Howell distinguishes about a dozen plant associations and discusses them under the headings: redwood forest, tanbark oak-madroño forest, oak-buckeye forest, Douglas fir forest, Bishop pine forest, chaparral, coastal brush, coastal grassland, hill and valley grassland, mountain meadow, serpentine grassland, vernal pools, streambank plants, lakeshore plants, freshwater marshes, salt marshes, dunes and beaches.

The major portion of the book is devoted to keys for identifying Marin County plants and to their enumeration. The nomenclature is commendably conservative. Useful notes and localities where a given species has been found are given under each. An appropriately abbreviated glossary and index, together with maps of the county and of Mt. Tamalpais, are appended to the main work. *Marin Flora* should prove to be particularly useful to botanists, amateur and professional alike, in the San Francisco Bay area.—R. C. ROLLINS.

¹ *MARIN FLORA: Manual of the Flowering Plants and Ferns of Marin County, California*, by JOHN THOMAS HOWELL. VII+ 323 pp. with 25 plates and 2 maps. \$4.50, University of California Press. 1949.

Volume 52, no. 615, including pages 49-60, was issued 17 March, 1950.

RATES FOR SPECIAL NUMBERS OF RHODORA

Many of the single numbers can be supplied only at special prices, as follows:

Vol. 12, no. 134: 50c	Vol. 37, no. 436: 70c	Vol. 46, no. 547: 50c
no. 138: 45c	no. 437: 50c	no. 548: 45c
Vol. 13, no. 151: 70c	no. 439: 60c	no. 550: 55c
Vol. 14, no. 163: 60c	no. 440: 60c	no. 551: 55c
Vol. 15, no. 171: 45c	no. 441: 50c	no. 552: 50c
Vol. 16, no. 182: 45c	no. 443: 55c	
Vol. 17, no. 193: 45c	no. 444: 55c	Vol. 47, no. 553: 75c
Vol. 18, no. 205: 50c	Vol. 38, no. 445: 50c	no. 554: 50c
Vol. 19, no. 224: 45c	no. 448: 70c	no. 555: 60c
no. 225: 50c	no. 450: 70c	no. 556: 75c
Vol. 21, no. 241: 45c	no. 455: 55c	no. 557: 75c
no. 243: 45c	no. 456: 50c	no. 558: 50c
Vol. 23, no. 265: 45c	Vol. 39, no. 458: 50c	no. 559: 75c
no. 268: 45c	no. 463: 55c	no. 560: 60c
no. 269: 45c	no. 464: 75c	no. 562: 85c
no. 270: 45c	no. 466: 55c	no. 563: 85c
no. 271: 45c	Vol. 40, no. 471: 55c	Vol. 48, no. 566: 60c
no. 274: 45c	no. 476: 50c	no. 567: 50c
no. 275: 45c	no. 477: 55c	no. 568: 60c
Vol. 24, no. 279: 45c	no. 478: 60c	no. 569: 50c
no. 283: 45c	no. 479: 55c	no. 570: 50c
Vol. 25, no. 296: 45c	Vol. 41, no. 482: 55c	no. 571: 60c
Vol. 26, no. 304: 50c	no. 486: 55c	no. 572: 50c
no. 305: 60c	no. 487: \$1	no. 573: 70c
no. 306: 45c	no. 488: 60c	no. 574: 70c
Vol. 28, no. 331: 45c	no. 489: 95c	no. 575: 70c
Vol. 29, no. 346: 45c	no. 490: 50c	no. 576: 50c
Vol. 30, no. 351: 50c	no. 491: 50c	
no. 356: 45c	Vol. 42, no. 499: 50c	Vol. 49, no. 577: 50c
no. 357: 45c	no. 500: 60c	no. 578: 60c
Vol. 31, no. 364: 50c	no. 502: 50c	no. 580: 60c
no. 369: 50c	no. 503: 70c	no. 581: 70c
no. 370: 50c	Vol. 43, no. 506: \$1	no. 582: 80c
Vol. 32, no. 374: \$1	no. 509: \$1	no. 583: 75c
no. 376: 45c	no. 512: 50c	no. 587: 50c
no. 382: 50c	no. 513: 50c	no. 588: 50c
no. 383: 45c	no. 514: 70c	
Vol. 33, no. 386: 60c	no. 515: 75c	Vol. 50, no. 589: 45c
no. 388: 45c	Vol. 44, no. 520: 70c	no. 590: 60c
no. 389: 45c	no. 525: 75c	no. 591: 40c
no. 391: \$1	no. 526: 75c	no. 592: 60c
Vol. 34, no. 403: 45c	no. 527: 70c	no. 593: 60c
no. 407: 45c	no. 528: 60c	no. 594: 40c
Vol. 35, no. 410: 50c	Vol. 45, no. 531: 60c	no. 595: 75c
no. 418: 50c	no. 532: 55c	no. 596: 85c
no. 419: 50c	no. 533: 55c	no. 597: 55c
Vol. 36, no. 425: 55c	no. 534: 75c	no. 598: 40c
no. 426: 50c	no. 535: 70c	no. 599: 60c
no. 429: 70c	no. 538: 85c	no. 600: 65c
no. 430: 55c	no. 539: 75c	
Vol. 37, no. 433: \$1	no. 540: 75c	Vol. 51, no. 603: 80c
no. 435: 60c	Vol. 46, no. 542: 50c	no. 604: 85c
	no. 544: 60c	no. 609: 75c
	no. 545: 55c	no. 610: 70c
	no. 546: 55c	no. 611: 70c
		no. 612: 70c

DUPLICATE BOOKS FOR SALE

- Braithwaite, R.** The British Moss-Flora. 3 vols. London. 1887-1905. 4°. 128 plates, ½ leather. **\$20.00**
- Dixon, H. N.** Student's Handbook of British Mosses; illustrations and keys by H. G. Jameson. London 1896. Plates 60. Cloth. **\$ 2.00**
- Same. Edition 2 revised with 65 plates. 1904. ½ leather. **\$ 3.50**
- Hitchcock, A. S. and Chase, A.** Grasses of the West Indies. (Contrib. U. S. Nat. Herb. 18, pt. 7, 1917). Covers worn. **\$ 2.00**
- Hitchcock, A. S. and Standley, P. C.** Flora of the District of Columbia and Vicinity. (Contrib. U. S. Nat. Herb. 21, 1919). **\$ 2.50**
- Maxon, W. R.** Studies of Tropical American Ferns—No. 3. (Contrib. U. S. Nat. Herb. 16, pt. 2, 1912). **\$ 2.00**
- Piper, C. V.** Flora of the State of Washington. (Contrib. U. S. Nat. Herb. 11, 1906). Worn covers and map lacking. **\$ 2.50**
- Safford, W. E.** Useful Plants of the Island of Guam. (Contrib. U. S. Nat. Herb. 9, 1905). Covers slightly worn. . . . **\$ 4.00**

Prices include cost of transportation in U. S. A.

Address Librarian,

GRAY HERBARIUM OF HARVARD UNIVERSITY
79 Garden St., Cambridge 38, Mass.

CARD-INDEX OF NEW GENERA, SPECIES AND VARIETIES OF AMERICAN PLANTS

For all students of American Plants the Gray Herbarium Card-index of Botanical Names is indispensable. It is a work of reference essential to scientific libraries and academies and all centers of botanical activity. It includes genera and species from 1885 to date. The subdivisions of species from 1885 to date are now included and from 1753 to 1886 are in the process of being inserted. Issued quarterly, at \$25.50 per thousand cards. Sets of paper facsimiles of issues 1-186 (through July, 1945) alphabeted in a single series can now be supplied for \$4,500 f.o.b. Massachusetts; issue 187 through 194 at \$22.50 per thousand.

GRAY HERBARIUM of Harvard University,
Cambridge 38, Mass., U. S. A.